

resources for enrichment, problem solving and contest preparation.

Grade 7

Scoring: There is *no penalty* for an incorrect answer. Each unanswered question is worth 2, to a maximum of 10 unanswered questions.



(D) 66 (E) 36





- Grade 7
- 18. Distinct points are placed on a circle. Each pair of points is joined with a line segment. An example with 4 points and 6 line segments is shown. If 6 distinct points are placed on a circle, how many line segments would there be?
 - (A) 13 (B) 16 (C) 30
 - (D) 15 (E) 14
- 19. If each of the four numbers 3, 4, 6, and 7 replaces a □, what is the largest possible sum of the fractions shown?
 - (A) $\frac{19}{12}$ (B) $\frac{13}{7}$ (C) $\frac{5}{2}$ (D) $\frac{15}{4}$ (E) $\frac{23}{6}$



20. Andy, Jen, Sally, Mike, and Tom are sitting in a row of five seats. Andy is not beside Jen. Sally is beside Mike. Who *cannot* be sitting in the middle seat?
(A) Andy (B) Jen (C) Sally (D) Mike (E) Tom

Part C: Each correct answer is worth 8.

- 21. A bicycle travels at a constant speed of 15 km/h. A bus starts 195 km behind the bicycle and catches up to the bicycle in 3 hours. What is the average speed of the bus in km/h?
 - (A) 65 (B) 80 (C) 70 (D) 60 (E) 50

22. In the *Coin Game*, you toss three coins at the same time. You win only if the 3 coins are all showing heads, or if the 3 coins are all showing tails. If you play the game once only, what is the probability of winning?

(A) $\frac{1}{6}$ (B) $\frac{1}{4}$ (C) $\frac{2}{27}$ (D) $\frac{2}{3}$ (E) $\frac{1}{3}$

23. Molly assigns every letter of the alphabet a *different* whole number value. She finds the value of a word by *multiplying* the values of its letters together. For example, if D has a value of 10, and I has a value of 8, then the word DID has a value of $10 \times 8 \times 10 = 800$. The table shows the value of some words. What is the value of the word MATH?

| Word | Value |
|------|-------|
| TOTE | 18 |
| TEAM | 168 |
| MOM | 49 |
| HOME | 70 |
| MATH | ? |

- (A) 19 (B) 840 (C) 420
- **(D)** 190 **(E)** 84
- 24. How many different pairs (m, n) can be formed using numbers from the list of integers $\{1, 2, 3, \ldots, 20\}$ such that m < n and m + n is even? (A) 55 (B) 90 (C) 140 (D) 110 (E) 50
- 25. Tanner wants to fill his swimming pool using two hoses, each of which sprays water at a constant rate. Hose A fills the pool in a hours when used by itself, where a is a positive integer. Hose B fills the pool in b hours when used by itself, where b is a positive integer. When used together, Hose A and Hose B fill the pool in 6 hours. How many different possible values are there for a? (A) 5 (B) 6 (C) 9 (D) 10 (E) 12